

A Propp-Wilson perfect simulation scheme for processes with long memory

Aurélien Garivier

June 30, 2011

Abstract

We describe a new “coupling from the past” algorithm for variable length Markov chains and random systems with perfect connections. This algorithm generalizes Propp and Wilson’s simulation scheme, and improves on existing algorithms in that it requires much less restrictive conditions on the kernel to converge. It proves interesting even in the simple case of finite order Markov chains in terms of speed of convergence. Chains of variable or infinite order are an old objet of consideration that raised considerable interest recently because of its applications in applied probability, from information theory to bio-informatics and linguistics.

Keywords: Perfect simulation, Context trees, Markov Chains of infinite order, Coupling from the past

1 Introduction

Since the seminal paper of [19], perfect simulation schemes for stationary Markov chains have been developped and applied in several fields of applied probabilities, from statistical physics to Bayesian statistics (see for example [16] and references therein, or [12] for a gentle introduction).

In 2002, Comets, Fernandez and Ferrari [5] have proposed an extension to processes with long memory: they provided a perfect simulation algorithm for stationary processes called random systems with complete connections [17, 18] or *chains of infinite order* [13]; these processes are characterized by a transition kernel which specifies, given an infinite sequence of past symbols, the probability distribution of the next symbol. The idea was, after [14, 15, 2], to exploite regenerative structures of these processes. Their algorithm relies on renewal properties, under a summable memory decay condition. As a by-product, the authors obtained the existence of stationary process for a given kernel, together with unicity properties under suitable hypotheses.

However, it appeared that these conditions on the kernel were quite restrictive, and actually not necessary. Foss [8] and Gallo [9] showed that different coupling schemes could be designed under alternative assumptions that do not even require the kernel to be continuous. Actually, the coupling scheme of Comets, Fernandez and Ferrari strongly relies on regeneration, contrary to Propp and Wilson’s algorithm: thus, it cannot compete with the latter on the simple case of Markov chains in terms of convergence time. Recently, De Santis and Piccioni tried to reconcile the two algorithms, by providing an hybrid methods

that works with two regimes: coalescence for short memory, and regeneration on long scales.

In this paper, we fill the gap between long and short scales, by providing a relatively elaborated and yet elegant coupling procedure that purely relies on coalescence. It generalizes Propp and Wilson's algorithm, in the sense that it is exactly similar for (first order) Markov chains. But it permits to handle arbitrary continuous infinite memory stationary processes, and even some non-continuous ones under particular conditions. The idea is to exploit *local* coalescence, instead of global loss of memory properties. We show that this perfect simulation scheme can be proved to converge under much less restrictive hypotheses than was known previously. Even when different schemes can be applied, it converges (possibly very significantly) faster. As they prove very useful in many applications (e.g. information Theory [20, 21] or bio-informatics [4]), we set a special focus on perfect simulation of finite, but large order Markov chains (or for Variable order Markov Chains, see [3]): our algorithm compares very favorably with Propp and Wilson's algorithm on the extended chain in terms of computational complexity.

The paper is organized as follows: Section 2 presents the required notation and definitions, together with a few first properties. Section 3 contains the conceptual description of the perfect simulation schemes. It relies on a coupling function that is described in great detail in Section 4. Section 5 contains the precise description of the algorithm. Then, Section 6 gathers elements of analysis of the complexity of the algorithm, while Section 7 illustrates the weakness of the assumptions required for the algorithm to converge.

2 Notation and background

2.1 Spaces and topologies

Following [5], we denote by G a finite alphabet. For $k \in \mathbb{N}$, we denote by G^{-k} the set of all sequences $(w_{-k}, \dots, w_{-1})$, and $G^* = \cup_{k \geq 0} G^{-k}$. By convention, ε denotes the empty sequence, and $G^0 = \{\varepsilon\}$. The set $G^{-\mathbb{N}^*}$ is the space of *histories*, and $w_{-\infty:-1} \in G^{-\mathbb{N}^*}$ will be denoted by $\underline{w}$. The history $\underline{w}$ such that for all $k < 0$, $w_k = 0$ will be denoted $\underline{0}$. For $w \in G^{-k}$, we note $|w| = k$ and for $\underline{w} \in G^{-\mathbb{N}^*}$, $|\underline{w}| = \infty$. For every non-positive integer n , we define the projection $\Pi^n : G^{-\mathbb{N}^*} \rightarrow G^n$ be defined by $\Pi_n(\underline{w}) = w_{n:-1}$.

Endowed with the ultrametric distance

$$\delta(\underline{w}, \underline{z}) = 2^{\sup\{k < 0 : w_k \neq z_k\}},$$

$G^{-\mathbb{N}^*}$ is a complete and compact set. A (closed and open) ball $B \subset G^{-\mathbb{N}^*}$ is a set $\{\underline{z}s : \underline{z} \in G^{-\mathbb{N}^*}\}$ for some $s \in G^*$: we denote $s = \mathcal{R}(B)$, and denote $B = \mathcal{T}(s)$. Observe that the radius of $\mathcal{T}(s)$ is $2^{-|s|}$. Note that $\mathcal{T}(\varepsilon) = G^{-\mathbb{N}^*}$. A mapping f defined on $G^{-\mathbb{N}^*}$ is called *piecewise constant* if there exists a family $\{s_j\}_{j \in \mathbb{N}}$ of elements of $G^{-\mathbb{N}^*}$ such that f is constant on each ball $\mathcal{T}(s_j)$.

For $-\infty \leq a \leq b \leq +\infty$ and $w \in G^{\mathbb{Z}}$, the sequence $(w_a, \dots, w_b)$ is denoted by $w_{a:b}$, and for two sequences $w_{a:b}$ and $z_{c:d}$, $-\infty \leq a \leq b < \infty$, $-\infty < c \leq d \leq \infty$, $w_{a:b}z_{c:d}$ denotes the concatenation of $w_{a:b}$ and $z_{c:d}$: $w_{a:b}z_{c:d} = (w_a, \dots, w_b, z_c, \dots, z_d)$. Remark: this notation is different from the convention taken in [5]. If $a > b$, $w_{a:b}$ is the empty sequence ε .

In the sequel, the set of probability distributions on G is denoted by $\mathcal{M}_1(G)$, and is endowed with the total variation (or L^1) distance

$$|p - q|_{TV} = \frac{1}{2} \sum_{a \in G} |p(a) - q(a)| = 1 - \sum_{a \in G} p(a) \wedge q(a) ,$$

where $x \wedge y$ denotes the minimum of x and y .

2.2 Probability transition kernels

A mapping $P : G^{-\mathbb{N}^*} \rightarrow \mathcal{M}_1(G)$ is called a *probability transition kernel*, and we denote the image of $\underline{w} \in G^{-\mathbb{N}^*}$ by $P(\cdot|\underline{w})$. We say that P is *continuous* if it is continuous from $(G^{-\mathbb{N}^*}, \delta)$ to $(\mathcal{M}_1(G), |\cdot|_{TV})$. For $s \in G^*$, we define the *oscillation* of P on the ball $\mathcal{T}(s)$ as:

$$\eta(s) = \sup \{ |P(\cdot|\underline{w}) - P(\cdot|\underline{z})|_{TV} : \underline{w}, \underline{z} \in \mathcal{T}(s) \} .$$

Lemma 1. *The following assertions hold:*

- (i) *P is continuous if and only if $\forall \underline{w} \in G^{-\mathbb{N}^*}, \eta(w_{-k:-1}) \rightarrow 0$ as k goes to infinity.*
- (ii) *P is continuous if and only if $\sup\{\eta(s) : s \in G^{-k}\} \rightarrow 0$ as k goes to infinity.*
- (iii) *P is uniformly continuous if and only if it is continuous.*

The two first assertions are trivial, and (iii) is a consequence of Heine's theorem on the compact set $G^{-\mathbb{N}^*}$.

We say that a (nonnecessarily stationary) process $(X_t)_{t \in \mathbb{Z}}$ with distribution ν on $G^{\mathbb{Z}}$ is *compatible* with kernel P if the latter is a version of the one-sided conditional probabilities of the former, that is:

$$\nu(X_i = g | X_{i+j} = w_j, j \in -\mathbb{N}^*) = P(g|\underline{w})$$

for all $i \in \mathbb{Z}, g \in G$ and ν -almost every $\underline{w}$. A classical but key remark is that $S_t = (\dots, X_{t-1}, X_t), t \in \mathbb{Z}$, is a Markov Chain on the compact ultrametric state space $G^{-\mathbb{N}^*}$, with transition kernel Q given by:

$$\forall \underline{w}, \underline{z} \in G^{-\mathbb{N}^*}, \quad Q(\underline{w}|\underline{z}) = P(w_{-1}|\underline{z}) \mathbb{1}_{w_{i-1}=z_i: i < 0} .$$

2.3 Coupling functions

A mapping $\phi : [0, 1[\times G^{-\mathbb{N}^*} \rightarrow G$ is called a *coupling* of P if, given a random variable U uniformly distributed on $[0, 1[$, $\phi(U, \underline{w})$ has distribution $P(\cdot|\underline{w})$ for all $\underline{w} \in G^{-\mathbb{N}^*}$.

In the sequel, we focus on couplings ϕ of P such that:

$$\forall s \in G^*, 0 \leq u < 1 - |G|\eta(s) \implies \phi(u, \cdot) \text{ is constant on } \mathcal{T}(s) . \quad (1)$$

The construction of such a coupling is detailed in Section 4.

Lemma 2. *If P is continuous and if ϕ is a coupling of P satisfying (1), then for all $u \in [0, 1[$ the mapping $\underline{w} \rightarrow \phi(u, \underline{w})$ is continuous, i.e. piecewise constant.*

Proof: let $u \in [0, 1[$; the uniform continuity of P implies that there exists ϵ such that if $\delta(\underline{w}, \underline{z}) \leq \epsilon$, then $|P(\cdot|\underline{w}) - P(\cdot|\underline{z})|_{TV} < (1 - u)/|G|$. But then Equation (1) implies that $\phi(u, \underline{w}) = \phi(u, \underline{z})$.

3 Description of the Perfect Simulation Scheme

Given a continuous transition kernel P , two questions arise:

1. does there exist a stationary distribution ν compatible with P ? In that case, is it unique?
2. if ν exists, how can we sample finite trajectories from that distribution?

More or less recently, [5, 7, 9] have contributed to answer these questions. Their approach is to show that there exists a simulation scheme drawing samples of ν , and this algorithm is based on the idea of coupling from the past. Following these authors, we address these questions by constructing a new perfect simulation scheme that converges faster, and more generally than its predecessors. We describe the general principle of this algorithm here.

Let n be a negative integer; in order to draw $(X_n, \dots, X_{-1})$ from a stationary distribution compatible with P , we use a semi-infinite sequence of i.i.d. random variables $(U_t)_{t < 0}$ on the probabilistic space $(\Omega, \mathcal{A}, P)$, each uniformly distributed on the interval $[0, 1]$. For each $t < 0$, let $f_t : G^{-\mathbb{N}^*} \rightarrow G^{-\mathbb{N}^*}$ be defined by $f_t(\underline{w}) = \underline{w}\phi(U_t, \underline{w})$; beware the index shift: if $\underline{z} = f_t(\underline{w})$ then $z_{-1} = \phi(U_t, \underline{w})$ and $z_i = w_{i+1}$ for $i < -1$.

For all integers $t < u$, let $F_{t:u} = f_{u-1} \circ \dots \circ f_t$, and let $F_t = F_{t:-1}$. For any negative integer n , let also $H_{t:u}^n = \Pi^n \circ F_{t:u}$ and let $H_t^n = H_{t:-1}^n$. As will be detailed below, for a continuous kernel Proposition 2 implies that $H_{t:u}^n$ is a piecewise constant mapping. Define

$$\tau(n) = \sup\{t < n : H_t^n \text{ is constant}\},$$

where by convention $\tau(n) = -\infty$ if for all $t < 0$, H_t^n is not constant. The sequence $(\tau(n))_n$ is a non-increasing sequence of stopping times with respect to the filtration $(\mathcal{F}_s)_s$, where $\mathcal{F}_s = \sigma(U_t : t \geq s)$. In [5, 7], conditions on P are given that ensure almost-sure finiteness for $\tau(n)$ (and bounded expectation). On the other hand, the authors prove that almost-sure finiteness for $\tau(n)$ is a sufficient condition to prove the existence and unicity of a stationary distribution ν compatible with P (see [5], Theorem 4.3 and corollaries 4.12 and 4.13). As a by-product, one obtains a simulation algorithm for sample paths of ν : if $U = (U_t)_{t \in \mathbb{Z}}$ is a sequence of independent, uniformly distributed random variables, one can define $\Phi : [0, 1]^{\mathbb{Z}} \rightarrow G^{\mathbb{Z}}$ such that $\Phi(U)_t = \phi(U_{t-1}, \Phi(U)_{-\infty:t-1})$ for all t , and

$$\nu = \mathbb{P}(\Phi(U) \in \cdot),$$

the law of $\Phi(U)$, is stationary and compatible with P .

But the conditions on P required in [5] are quite restrictive: with the notation of Section 4 below, it is assumed that

$$\sum_{m \geq 0} \prod_{k=0}^m A_k^- < \infty.$$

This condition require in particular that $A_1^- = \sum_{g \in G} \inf \{P(g|\underline{z}) : \underline{z} \in G^{-\mathbb{N}^*}\} > 0$. The authors prove that, under these contions, the process regenerates, and that the stopping time

$$\tau'(n) = \sup\{t < n : H_t^t \text{ is constant}\}$$

is almost-surely finite, using a Kalikow-type decomposition of the kernel P as a mixture of Markov chains of all orders. . This is obviously sufficient to ensure that $\tau(n)$ is finite, but this not necessary by far. Consider the example of order 1 Markov chains: while Propp and Wilson [19] have shown that $\tau(n)$ is almost surely finite for every mixing chain (and, besides, that $\tau(n)$ has the same order of magnitude as the mixing time of the chain), $\tau'(n)$ is easily seen to be almost surely infinite if the Dobrushin coefficient A_1^- of the chain is 0. The contribution of this paper is precisely to fill the gap, by providing for general continuous kernels a Propp-Wilson procedure that may converge even if the process is not regenerating. In fact, for Markov chains of order 1, $\phi(u, \underline{w})$ depends only on w_{-1} and the algorithm presented in this paper is simply Propp and Wilson's exact sampling procedure.

Since the publication of [5], [9, 7] have generalized these results, showing that they are not necessary, by proposing other particular conditions covering different cases. Gallo [9] shows that P needs not be continuous to ensure the existence of ν , nor to ensure the finiteness of $\tau(n)$: he gives an example of a non-continuous regenerating chain, see also Example 7.2 below. In [7], De Santis and Piccioni propose another algorithm wich mixes the ideas of [5] and [19]: they propose an hybrid simulation scheme working with a Markov regime and a long-memory regime. Our approach is different and more general: we describe a single procedure that generalizes the sampling schemes of [5] and [19] in a single, unified framework.

4 Construction of the coupling function

Provide G with any order, so that $G^{-\mathbb{N}^*}$ can be equiped with the lexicographic order : $\underline{w} < \underline{z}$ if there exists $k \in -\mathbb{N}$ such that $\forall j > k, w_j = z_j$ and $w_k < z_k$. In this section and in the following, we consider a fixed continuous kernel P . The continuity of P is locally characterized by some coupling factors that we define here together with quantities that will prove useful for the construction of the coupling function ϕ : for all $g \in G$, let $A_{-1}(\varepsilon) = a_{-1}(g|\varepsilon) = 0$ and for $k \in \mathbb{N}, w \in G^*$, let

$$\begin{aligned} a_k(g|w_{-k:-1}) &= \inf\{P(g|\underline{z}) : \underline{z} \in \mathcal{T}(w_{-k:-1})\} , \\ A_k(w_{-k:-1}) &= \sum_{g \in G} a_k(g|w_{-k:-1}) , \\ A_k^- &= \inf_{s \in G^{-k}} A_k(s) , \\ \alpha_k(g|w_{-k:-1}) &= A_{k-1}(w_{-k+1:-1}) + \sum_{h < g} \{a_k(h|w_{-k:-1}) - a_{k-1}(h|w_{-k+1:-1})\} , \\ \beta_k(g|w_{-k:-1}) &= A_{k-1}(w_{-k+1:-1}) + \sum_{h \leq g} \{a_k(h|w_{-k:-1}) - a_{k-1}(h|w_{-k+1:-1})\} . \end{aligned}$$

Note that, with our conventions, $a_0(g|\varepsilon) = \inf\{P(g|\underline{z}) : \underline{z} \in G^{-\mathbb{N}^*}\}$. Moreover, if $s, s' \in G^*$ are such that $s \succeq s'$, then for all $g \in G$ it holds that $a_k(g|s) \geq a_k(g|s')$, $A_k(s) \geq A_k(s')$ and the sequence $(A_k^-)_k$ is increasing.

Proposition 1. *The coupling factors of the kernel P satisfy the following inequalities: for all $s \in G^*$,*

$$1 - |G|\eta(s) \leq A_{|s|}(s) \leq 1 - \eta(s) . \quad (2)$$

Proof. For the upper-bound, observe that

$$\begin{aligned} \eta(s) &= \sup \{ |P(\cdot|\underline{w}) - P(\cdot|\underline{z})|_{TV} : \underline{w}, \underline{z} \in \mathcal{T}(s) \} \\ &= \sup \left\{ 1 - \sum_{a \in G} P(a|\underline{w}) \wedge P(a|\underline{z}) : \underline{w}, \underline{z} \in \mathcal{T}(s) \right\} \\ &= 1 - \inf \left\{ \sum_{a \in G} P(a|\underline{w}) \wedge P(a|\underline{z}) : \underline{w}, \underline{z} \in \mathcal{T}(s) \right\} \\ &\leq 1 - \sum_{a \in G} \inf \{ P(a|\underline{w}) \wedge P(a|\underline{z}) : \underline{w}, \underline{z} \in \mathcal{T}(s) \} \\ &= 1 - \sum_{a \in G} \inf \{ P(a|\underline{w}) : \underline{w} \in \mathcal{T}(s) \} \\ &= 1 - A_{|s|}(s) . \end{aligned}$$

For the lower-bound, observe that for all $\underline{z} \in \mathcal{T}(s)$, as $|P(\cdot|\underline{0}s) - P(\cdot|\underline{z})|_{TV} \leq \eta(s)$, it holds that $P(a|\underline{z}) \geq P(a|\underline{0}s) - \eta(s)$ for all $a \in G$ and thus

$$\begin{aligned} A_{|s|}(s) &= \sum_{a \in G} \inf \{ P(a|\underline{z}) : \underline{z} \in \mathcal{T}(s) \} \\ &\geq \sum_{a \in G} P(a|\underline{0}s) - \eta(s) \\ &= 1 - |G|\eta(s) . \end{aligned}$$

□

There is a minor improvement for this result: one can show that

$$A_{|s|}(s) = \sum_{a \in G} \inf \{ P(a|z) : \underline{z} \in \mathcal{T}(s) \} \geq 1 - (|G| - 1)\eta(s) .$$

This bound cannot be improved in general.

Proposition 2. *The three assertions are equivalent:*

- (i) *the kernel P is continuous;*
- (ii) $\forall \underline{w} \in G^{-\mathbb{N}^*}, A_k(w_{-k:-1}) \rightarrow 1$ as $k \rightarrow \infty$;
- (iii) $A_k^- \rightarrow 1$ as k goes to infinity.

Proof: The equivalence between (i) and (ii) is a simple consequence of Lemma 1 and Proposition 1. Similarly, (iii) follows from (i): if P is continuous on the compact set $G^{-\mathbb{N}^*}$, then it is uniformly continuous, and

$$f(k) = \sup_{s \in G^{-k}} \eta(s) \rightarrow 0$$

as k goes to infinity. But by Proposition 1, $A_k^- \geq 1 - |G|f(k)$. Finally, (iii) implies (ii).

The equivalence of (ii) and (iii) can also be proved as a consequence of Dini's first theorem: defining $\tilde{A}_k(\underline{w}) = A_k(w_{-k:-1})$, the sequence $(\tilde{A}_k)_k$ is an increasing sequence of continuous functions simply converging to the (continuous) constant function 1, thus the convergence is uniform.

Proposition 3. *For every $\underline{w} \in G^{-\mathbb{N}^*}$,*

$$[0, 1[= \bigcup_{g \in G, k \in \mathbb{N}} [\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[.$$

In other words: let $u \in [0, 1[$. For all $\underline{w} \in G^{-\mathbb{N}^*}$, there exists a unique $k \in \mathbb{N}$ and a unique $g \in G$ such that $u \in [\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[$.

Proof: Let m (resp. M) be the minimal (resp. maximal) element of G . Then $\alpha_k(m|w_{-k:-1}) = A_{k-1}(w_{-k+1:-1})$, $\beta_k(M|w_{-k:-1}) = A_k(w_{-k:-1})$, and

$$[A_{k-1}(w_{-k+1:-1}), A_k(w_{-k:-1})[= \bigcup_{g \in G} [\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[.$$

The result follows from the continuity assumption: $A_{-1}(\varepsilon) = 0$ and $A_k(w_{-k:-1}) \rightarrow 1$ as k goes to infinity. Figure 1 illustrates Proposition 3 on a three-symbols alphabet.




Figure 1: Graphical representation of a coupling ϕ on alphabet $\{0, 1, 2\}$: for each $w_{-k:-1}$, the intervals $[\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[$ are represented in blue ($g = 0$), red ($g = 1$) and green ($g = 2$). For example, $P(1|1) = \alpha_0(1|\varepsilon) + \alpha_1(1|1) = 1/8 + 1/4$, and $P(0|00) = \alpha_0(0|\varepsilon) + \alpha_1(0|0) + \alpha_2(0|00) = 1/4 + 1/8 + 0$.

Proposition 3 allows the following definition:

Definition 1. *The mapping $\phi : [0, 1[\times G^{-\mathbb{N}^*} \rightarrow G$ is defined as follows:*

$$\phi(u, \underline{w}) = \sum_{g \in G, k \in \mathbb{N}} g \mathbb{1}_{[\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[}(u).$$

In words, for every $u \in [0, 1[$ and for every $\underline{w} \in G^{-\mathbb{N}^*}$, $\phi(u, \underline{w})$ is the unique symbol $g \in G$ such that there exists $k \in \mathbb{N}$ satisfying: $u \in [\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})[$.

Proposition 4. ϕ is a coupling function satisfying Equation (1):

$$\forall s \in G^*, \forall u \in [0, 1], \forall \underline{w}, \underline{z} \in \mathcal{T}(s), \quad u < A_{|s|}(s) \implies \phi(u, \underline{w}) = \phi(u, \underline{z}) .$$

For every $s \in G^*$ and every $u < A_{|s|}(s)$ we denote in the sequel $\phi(u, s) = \phi(u, \underline{w})$ for all $\underline{w} \in \mathcal{T}(s)$.

Proof: We need to prove that if $U \sim \mathcal{U}([0, 1])$, then for all $\underline{w} \in G^{-\mathbb{N}^*}$ then random variable $\phi(U, \underline{w})$ has distribution $P(\cdot | \underline{w})$. It is sufficient to prove that for all $g \in G$,

$$\sum_{l=0}^{\infty} \beta_k(g|w_{-l:-1}) - \alpha_k(g|w_{-l:-1}) = P(g|\underline{w}).$$

Actually, it holds that

$$\sum_{l=0}^k \beta_k(g|w_{-l:-1}) - \alpha_k(g|w_{-l:-1}) = \sum_{l=0}^k a_l(g|w_{-l:-1}) - a_{l-1}(g|w_{-l+1:-1}) = a_k(g|w_{-l:-1}).$$

As an increasing sequence upper-bounded by $P(g|\underline{w})$, $a_k(g|w_{-k:-1})$ has a limit $Q(g|\underline{w}) \leq P(g|\underline{w})$ as k tends to infinity. By continuity,

$$\sum_{g \in G} Q(g|\underline{w}) = \sum_{g \in G} \lim_{k \rightarrow \infty} a_k(g|w_{-k:-1}) = \lim_{k \rightarrow \infty} \sum_{g \in G} a_k(g|w_{-k:-1}) = \lim_{k \rightarrow \infty} A_k(w_{-k:-1}) = 1,$$

and as $\sum_{g \in G} P(g|\underline{w}) = 1$ this implies that for all $g \in G$, $Q(g|\underline{w}) = P(g|\underline{w})$.

The last part of the proposition is immediate: for $\underline{w}, \underline{z} \in \mathcal{T}(s)$, $\forall k \leq |s|$, $w_{-k:-1} = z_{-k:-1}$ and

$$\bigsqcup_{g \in G, k \leq |s|} [\alpha_k(g|w_{-k:-1}), \beta_k(g|w_{-k:-1})] = \bigsqcup_{g \in G, k \leq |s|} [\alpha_k(g|z_{-k:-1}), \beta_k(g|z_{-k:-1})] = [0, A_{|s|}(s)] .$$

Having defined the coupling function, we can now describe the simulation scheme outlined in Section 3.

5 The Coupling from the Past Algorithm

5.1 Complete suffix Dictionaries

Let $h \in G^* \cup G^{-\mathbb{N}^*}$. If $s \in G^*$ is such that $|h| \geq |s|$ and $h_{-|s|:-1} = s$, we say that s is a *suffix* of h and we write $h \succeq s$.

A (finite or infinite) set of words $D \subset \mathcal{P}(G^*)$ is called a *complete suffix dictionary* (CSD) if one of the following equivalent properties is satisfied:

- every sequence $\underline{w} \in G^{-\mathbb{N}^*}$ has a unique suffix in D :

$$\forall \underline{w} \in G^{-\mathbb{N}^*}, \exists ! s \in D : \underline{w} \succeq s ;$$

- $\{\mathcal{T}(s) : s \in D\}$ is a partition of $G^{-\mathbb{N}^*}$:

$$G^{-\mathbb{N}^*} = \bigsqcup_{s \in D} \mathcal{T}(s) .$$

The CSD D can be represented by a *trie*, that is, a tree with edges labelled by elements of G such that the path from the root to any leaf is labelled by an element of D , see Figure 2. Accordingly, we define the *depth* of D as the depth of this trie:

$$d(D) = \sup\{|s| : s \in D\}$$

The smallest possible CSD is $\{\epsilon\}$ (its trie is reduced to the root): it has depth 0 and size 1. The second smallest is G , it has depth 1. If D and D' are two CSD such that $\forall s \in D', s \succeq D$, then we note $D' \succeq D$. This means that the trie representing D' entirely shadows that of D .




Figure 2: Left: the trie representing the Complete Suffix Dictionary $D = \{00, 01, 11\}$. Right: $\{00, 10, 001, 101, 11\} \succeq \{0, 01, 11\}$. Both examples concern the binary alphabet.

To every CSD D we associate a mapping, denoted by $\vec{D}$, defined on $G^* \cup G^{-\mathbb{N}^*}$ as follows:

- if there exists $s \in D$ such that $h \succeq s$, we note $h \succeq D$ and we define $\vec{D}(h) = \{s\}$;
- otherwise, we define $\vec{D}(h) = \{s \in D : s \succeq h\}$.

5.2 Piecewise constant functions and Complete Suffix Dictionaries

Definition 2. For a CSD D , we say that a function f defined on $G^{-\mathbb{N}^*}$ is D -constant if

$$\forall s \in D, \forall w \in \mathcal{T}(s), f(\underline{w}) = f(\underline{s}).$$

Then, for every $h \in G^{-\mathbb{N}^*} \cup G^*$ we define $f(h) = f(\mathcal{T}(h)) = f(\vec{D}(h))$ and note that if $h \succeq D$, $f(h)$ is a singleton.

Definition 3. A function f that is piecewise constant admits a minimal CSD D^f , which is defined as the CSD with minimal cardinality such that f is constant on each of its elements.

The minimal CSD D^f is such that if $s \in D^f$, there exists $w \in G^*$ such that $s' = ws_{-|s|+1:-1} \in D$ and $f(s) \neq f(s')$. If f is D -constant, then D^f can be obtained by recursive pruning of D : as long as it is possible, prune the nodes

whoses children are leaves with the same value for f . Pruning is illustrated at the right side of Figure 3.

We now turn back to the continuous kernel P and to its coupling function ϕ . For each $u \in [0, 1[$, Proposition 2 shows that the mapping $\phi(u, \cdot)$ is piecewise constant; we denote by $D(u) = D^{\phi(u, \cdot)}$ its minimal CSD. In the sequel, we show how the mappings H_t^n defined in Section 3 can be recursively constructed.

5.3 Recursive construction of H_t^n

Let t and n be negative integers. The mapping H_t^n being piecewise constant, we can define $D_t^n = D^{H_t^n}$.

First, observe that for $t \geq n$ and for $n \leq k \leq t$, it holds that $(H_t^n(\underline{w}))_k = w_k$, and H_t^n cannot be constant: $\tau(n) < n$. Besides, for $t \geq n$ it is easy to deduce H_t^t from H_t^{t+1} . Thus, the coupling algorithm successively considers $H_{-1}^{-1}, H_{-2}^{-2}, \dots, H_n^n, H_{n-1}^{n-1}, H_{n-2}^{n-2}, \dots$, and stops for the largest value of $t < n$ such that H_t^n is constant. We propose the following recursive construction:

- Initialization: let $D_{-1}^1 = G$ and $\forall g \in G, \forall \underline{w} \in \mathcal{T}(s), H_{-1}^1(\underline{w}) = g$.
- For $t < -1$, let for each $s \in D(U_t)$ denote $\{g_t(s)\} = \phi(U_t, s)$, and define $E_t^n(s)$ as follows:
 - if $sg_t(s) \succeq D_{t+1}^n$, let $E_t^n(s) = \{s\}$;
 - otherwise, let

$$E_t^n(s) = \bigcup_{hg_t(s) \in D_{t+1}^n(sg_t(s))} \{h\}.$$

- Let

$$E_t^n = \bigcup_{s \in D(U_t)} E_t^n(s).$$

E_t^n is a CSD, and H_t^n is E_t^n -constant.

- D_t^n is obtained by pruning E_t^n ; besides, for $t = n$, D_t^t is equal to D_t^{t+1} unless $D_t^{t+1} = \{\epsilon\}$, in which case $D_t^t = G$.

This recursion is illustrated in Figure 3.

Proof: Let us first check that E_t^n is a CSD. Note that every $h \in E_t^n(s)$ is such that $h \succeq s$; let $\underline{w} \in \mathcal{T}(s)$, and let $\{hg_t^s\} = D_{t+1:n}(\underline{w}g_n^s)$. Then $h \in E_t^n(s)$ and $\underline{w} \in \mathcal{T}(h)$, so that $\mathcal{T}(s) = \sqcup_{h \in E_t^n(s)} \mathcal{T}(h)$. The result follows, as $G^{-\mathbb{N}^*} = \sqcup_{s \in D(U_t)} \mathcal{T}(s)$.

Now, we can prove that H_t^n is E_t^n -constant. We proceed by induction on $t < 0$. The case $t = -1$ is clear. For $t < -1$, let $h \in E_t^n$, then by construction $h \succeq D(U_t)$, and thus $\phi(U_t, h)$ is the singleton $\{g_t(s)\}$ for $s = D(U_t)(h)$. Now, observe that $hg_t(s) \succeq D_{t+1}^n$, thus $H_t^n(h) = H_{t+1}^n(hg_t(s))$, which is a singleton by the induction hypothesis.

Besides, for $t = n$, first assume that $D_t^{t+1} \neq \{\epsilon\}$; then $|h| \geq 1$ and $H_t^t(h) = h_{-1}H_{t+1}^{t+1}(hg_t(s))$ is a singleton, so that H_t^t is D_t^{t+1} -constant. Otherwise, if $D_t^{t+1} = \{\epsilon\}$, H_t^{t+1} is constant equal to, say, $s' \in G^{t+1}$, then $\forall \underline{w} \in G^{-\mathbb{N}^*}, \forall g \in G, H_t^t(\underline{w}g) = gs'$, and thus $D(H_t^t) = G$.

Having described the coupling algorithm, we now give some elements of analysis concerning its complexity. We first give sufficient conditions for the algorithm to terminate, and then we focus on the special but interesting case of bounded memory processes.




Figure 3: Obtaining D_t^n from D_t and D_{t+1}^n . For each function $\phi(U_t, \cdot)$, D_{t+1}^n and D_t^n , we represent a CSD on which it is constant, and the values taken in each leaf; here, $G = \{0, 1\}$.

6 Bounding the size of D_t^n

6.1 Almost sure termination of the coupling scheme

In general, the CSD D_t^n can be arbitrary large with positive probability. In [5], conditions are given that ensure the finiteness of $\tau(n)$ defined above, from which bounds on D_t^n can be deduced. However, these conditions are quite restrictive: in particular, it is necessary that $A_0(\varepsilon) > 0$. [7] somewhat release these conditions using an hybrid simulation scheme, allowing for $A_0(\varepsilon) = 0$.

A very crude bound (ignoring the 'Propp-Wilson' coalescence possibilities of the algorithm) in the following: denoting $L_t^n = d(D_t^n)$, an immediate inspection of the recurrence 5.3 leads to:

$$\begin{cases} L_t^n \leq \max\{X_t, L_{t+1}^n - 1\} & \text{if } t < n, \text{ and} \\ L_t^n \leq \max\{X_t, L_{t+1}^{t+1} - 1, 1\} & \text{if } t \geq n. \end{cases}$$

where the $X_t = d(D(U_t))$ are i.i.d. random variables such that $\forall k \in \mathbb{N}, \mathbb{P}(X_t \leq k) = A_k^-$. Thus,

$$\mathbb{P}(L_t^n \leq k) \geq \mathbb{P}(L_{t+1}^n \leq k+1) A_k^- \geq \prod_{j=k}^{k-t-1} A_j^-.$$

This bound permits to recover the results of [5], and one obtains:

$$\mathbb{E}[L_t^n] \leq \sum_{k=1}^n 1 - \prod_{j=k}^{\infty} A_j^-.$$

6.2 The case of Finite Context Trees

There is at least one case where it is easy to upper-bound the size of $\mathcal{S}_t$ independently of $t \leq n$: when the kernel P actually defines a *Finite Context Tree*, that is, when the mapping $\underline{w} \rightarrow P(\cdot|\underline{w})$ is piecewise constant – then, denoting by D the minimal CSD of this mapping, it holds that $\forall \underline{w} \in G^{-\mathbb{N}^*}, P(\cdot|\tilde{D}(\underline{w})) = \{P(\cdot|\underline{w})\}$. Even in that case, the simulation scheme described above is interesting: although the “plain” Propp-Wilson algorithm could be applied on the first order Markov chain $(X_{t+1:t+d(D)})_{t \in \mathbb{Z}}$ on the extended state space $G^{d(D)}$, the computational complexity of such an algorithm might well become rapidly intractable if the depth $d(D)$ is not very small, whereas the following property shows that our algorithm keeps a possibly much more limited complexity. Such qualities of *parcimony* are precisely the reason why finite context trees have proved successful in many applications, from Information Theory and universal coding (see [20, 21, 6, 11]) to biology ([1, 4]) or linguistics [10].

We say that a CSD D is *prefix-closed* if every prefix of any sequence in D is the suffix of an element of D :

$$\forall s \in D, \forall k \leq |s|, \exists w \in D : w \succeq s_{-|s|:-k} .$$

A prefix-closed CSD satisfies the following property:

Lemma 3. *If D is a prefix-closed CSD, then for all $h \in D$ (or, equivalently, for all $h \succeq D$) and for all $a \in G$, $ha \succeq D$.*

Proof: If $h \in G^*$ is such that for some $a \in G$, $ha \succeq D$ does not hold, then (as D is a CSD) there exists $s \in D$ and $s' \in G^* \setminus \{\epsilon\}$ such that $s = s'ha$. But then $s'h$ is a prefix of s and, by the prefix-closure property, there exists $w \in D$ such that $w \succeq s'h$. Thus, one cannot have $h \succeq D$.

We define the *prefix closure* $\overleftarrow{D}$ of a CSD D as the minimal prefix-closed set containing D , that is, the set of maximal elements (for the partial order $\succeq$) of

$$\tilde{D} = \{s_{-|s|:-k} : s \in D, k \leq |s|\} .$$

In other words, $\overleftarrow{D}$ is the smallest set such that for all $w \in \tilde{D}$ there exists $s \in \overleftarrow{D}$ such that $s \succeq w$.

Proposition 5. *For any complete suffix dictionary D , $\overleftarrow{D}$ is a complete suffix dictionary such that $|\overleftarrow{D}| \leq |D| \times d(D)$.*

The majoration is trivial: $|\overleftarrow{D}| \leq |\tilde{D}| \leq |D| \times d(D)$. It is extremely pessimistic in general: many CSD are already prefix-closed, and for most CSD $|\overleftarrow{D}|$ is of the same order of magnitude as $|D|$. But in fact, for each positive integer n , there exists a CSD D of size n such that $|\overleftarrow{D}| \geq c|D|^2$ for some constant $c \approx 0.4$.

Now, assume that $D \neq \{\epsilon\}$, i.e. that P is not memoryless.

Proposition 6. *For each $t \leq n$, $\forall k < t$, $\overleftarrow{D} \succeq D_t^n$. Thus, a fortiori, $|D_t^n| \leq |\overleftarrow{D}| \leq |D| \times d(D)$.*

Proof: We show that $\overleftarrow{D} \succeq D_t^n$ by induction on t . First, as P is not memoryless, $\overleftarrow{D} \succeq D_{-1}^{-1} = G$. Second, assume that $\overleftarrow{D} \succeq D_{t+1}^n$: it is sufficient to prove




Figure 5: Convergence of the simulation scheme

three times a geometric variable of parameter $3/(2\sqrt{2})-1$. To simplify notations, denote $H_t = H_t^{-1}$. For every $t < -2$ if $U_{t-1} \leq 1 - 1/\sqrt{2}$, if $U_t > 1 - 1/\sqrt{2}$ and if $U_{t+1} \leq 1 - 1/\sqrt{2}$ then for every $\underline{w} \in G^{-\mathbb{N}^*}$ we have:

- $U_{t+1} \leq 1 - 1/\sqrt{2}$ implies $f_{t+1}(\underline{w}1) = \underline{w}10$ and $H_{t+1}(\underline{w}1) = H_{t+2}(\underline{0})$;
- $U_t > 1 - 1/\sqrt{2}$ implies $f_t(\underline{w}01) = \underline{w}011$ and $H_t(\underline{w}01) = H_{t+1}(\underline{w}011) = H_{t+2}(\underline{0})$ on the other hand, $f_t(\underline{w}0) = \underline{w}01$ and $H_t(\underline{w}0) = H_{t+1}(\underline{w}01) = H_{t+2}(\underline{0})$;
- $U_{t-1} \leq 1 - 1/\sqrt{2}$ implies $f_{t-1}(\underline{w}1) = \underline{w}10$ and $f_{t-1}(\underline{w}0) = \underline{w}01$, so that $H_{t-1}(\underline{w}0) = H_{t-1}(\underline{w}1) = H_{t+2}(\underline{0})$, and $\tau \geq t - 1$.

For every negative integer k let $E_k = \{U_{3k-1} \leq 1 - 1/\sqrt{2}\} \cap \{U_{3k} > 1 - 1/\sqrt{2}\} \cap \{U_{3k+1} \leq 1 - 1/\sqrt{2}\}$. The $(E_k)_{k < 0}$ are independent events of probability $3/(2\sqrt{2}) - 1$, which gives the result.

Thus, the algorithm converges quickly. However, the dictionaries involved in the simulation can be very large: in fact, it is easy to see that the depth X_t of $D(U_t)$ has no expectation: $\mathbb{P}(X_t \geq k) = 1/\sqrt{k}$. Of course, as D_t^n has a very special shape, ad hoc modifications of the algorithm would easily allow, here, to draw arbitrary long samples with low computational complexity.

7.2 A non-continuous process

Actually, a simple modification of this example show that continuity is absolutely not necessary to ensure convergence : the proof also applies to any kernel P' such that for $0 \leq k \leq 2$, $P'(0|01^k) = 1 - 1/\sqrt{k}$, and for any $\underline{w} \in G^{-\mathbb{N}^*}$, $P'(0|\underline{w}11) \geq 1 - 1/\sqrt{2}$. Such a phenomenon has been studied in [9]: Gallo gives sufficient conditions on the shape of the trees, together with bounds on transition probabilities, that ensure convergence of his coupling scheme. However, his approach is quite different and does not cover the examples presented here.

Acknowledgments

The authors thanks Sandro Gallo, Antonio Galves and Florencia Leonardi (Numecc, Sao Paulo) for stimulating and enlightening discussions on chains of infinite memory. This work was supported by USP-COFECUB (grant 2009.1.820.45.8).

References

- [1] G. Bejerano and G. Yona. Variations on probabilistic suffix trees: statistical modeling and prediction of protein families. *Bioinformatics*, 17(1):23–43, 2001.
- [2] Henry Berbee. Chains with infinite connections: uniqueness and Markov representation. *Probab. Theory Related Fields*, 76(2):243–253, 1987.
- [3] P. Bühlmann and A. J. Wyner. Variable length Markov chains. *Ann. Statist.*, 27:480–513, 1999.
- [4] J. R. Busch, P. A. Ferrari, A. G. Flesia, R. Fraiman, S. P. Grynberg, and F. Leonardi. Testing statistical hypothesis on random trees and applications to the protein classification problem. *Annals of applied statistics*, 3(2), 2009.
- [5] Francis Comets, Roberto Fernández, and Pablo A. Ferrari. Processes with long memory: regenerative construction and perfect simulation. *Ann. Appl. Probab.*, 12(3):921–943, 2002.
- [6] I. Csiszár and Z. Talata. Context tree estimation for not necessarily finite memory processes, via BIC and MDL. *IEEE Trans. Inform. Theory*, 52(3):1007–1016, 2006.
- [7] Mauro Piccioni Emilio De Santis. A general framework for perfect simulation of long memory processes, April 2010.
- [8] S. G. Foss, R. L. Tweedie, and J. N. Corcoran. Simulating the invariant measures of Markov chains using backward coupling at regeneration times. *Probab. Engrg. Inform. Sci.*, 12(3):303–320, 1998.
- [9] Sandro Gallo. Chains with unbounded variable length memory: perfect simulation and visible regeneration scheme, 2010.
- [10] A. Galves, C. Galves, J. Garcia, N.L. Garcia, and F. Leonardi. Context tree selection and linguistic rhythm retrieval from written texts. *ArXiv: 0902.3619*, pages 1–25, 2010.
- [11] A. Garivier. Consistency of the unlimited BIC context tree estimator. *IEEE Trans. Inform. Theory*, 52(10):4630–4635, 2006.
- [12] Olle Häggström. *Finite Markov chains and algorithmic applications*, volume 52 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2002.
- [13] Theodore E. Harris. On chains of infinite order. *Pacific J. Math.*, 5:707–724, 1955.

- [14] S. P. Lalley. Regenerative representation for one-dimensional Gibbs states. *Ann. Probab.*, 14(4):1262–1271, 1986.
- [15] S. P. Lalley. Regeneration in one-dimensional Gibbs states and chains with complete connections. *Resenhas*, 4(3):249–281, 2000.
- [16] D. J. Murdoch and P. J. Green. Exact sampling from a continuous state space. *Scand. J. Statist.*, 25(3):483–502, 1998.
- [17] Octav Onicescu and Gheorghe Mihoc. Sur les chaînes de variables statistiques. *Bull. Sci. Math.*, 59:174–192, 1935.
- [18] Octav Onicescu and Gheorghe Mihoc. Sur les chaînes statistiques. *Comptes Rendus de l'Académie des Sciences de Paris*, 200:511–512, 1935.
- [19] James Gary Propp and David Bruce Wilson. Exact sampling with coupled Markov chains and applications to statistical mechanics. In *Proceedings of the Seventh International Conference on Random Structures and Algorithms (Atlanta, GA, 1995)*, volume 9, pages 223–252, 1996.
- [20] J. Rissanen. A universal data compression system. *IEEE Trans. Inform. Theory*, 29(5):656–664, 1983.
- [21] F.M.J. Willems, Y.M. Shtarkov, and T.J. Tjalkens. The context-tree weighting method: Basic properties. *IEEE Trans. Inf. Theory*, 41(3):653–664, 1995.